

CONSTRUCTING UNIVERSALLY SMALL SUBSETS OF A GIVEN PACKING INDEX IN POLISH GROUPS

TARAS BANAKH AND NADYA LYASKOVSKA

ABSTRACT. A subset of a Polish space X is called *universally small* if it belongs to each ccc σ -ideal with Borel base on X . Under CH in each uncountable Abelian Polish group G we construct a universally small subset $A_0 \subset G$ such that $|A_0 \cap gA_0| = \mathfrak{c}$ for each $g \in G$. For each cardinal number $\kappa \in [5, \mathfrak{c}^+]$ the set A_0 contains a universally small subset A of G with sharp packing index $\text{pack}^\sharp(A_\kappa) = \sup\{|\mathcal{D}|^+ : \mathcal{D} \subset \{gA\}_{g \in G} \text{ is disjoint}\}$ equal to κ .

1. INTRODUCTION

This paper is motivated by a problem of Dikranjan and Protasov [4] who asked if the group of integers $\mathbb{Z}$ contains a subset $A \subset \mathbb{Z}$ such that the family of shifts $\{x + A\}_{x \in \mathbb{Z}}$ contains a disjoint subfamily of arbitrarily large finite cardinality but does not contain an infinite disjoint subfamily. This problem can be reformulated in the language of packing indices $\text{pack}(A)$ and $\text{pack}^\sharp(A)$, defined for any subset A of an group G by the formulas:

$$\text{pack}(A) = \sup\{|\mathcal{D}| : \mathcal{D} \subset \{gA\}_{g \in G} \text{ is a disjoint subfamily}\}$$

and

$$\text{pack}^\sharp(A) = \sup\{|\mathcal{D}|^+ : \mathcal{D} \subset \{gA\}_{g \in G} \text{ is a disjoint subfamily}\}.$$

So, actually Dikranjan and Protasov asked about the existence of a subset $A \subset \mathbb{Z}$ with packing index $\text{pack}^\sharp(A) = \aleph_0$. This problem was answered affirmatively in [1] and [2]. Moreover, in [7] the second author proved that for any cardinal κ with $2 \leq \kappa \leq |G|^+$ and $\kappa \notin \{3, 4\}$ in any Abelian group G there is a subset $A \subset G$ with packing index $\text{pack}^\sharp(A) = \kappa$. By Theorem 6.3 of [3], such a set A can be found in any subset $L \subset A$ with Packing index $\text{Pack}(L) = 1$ where

$$\text{Pack}(A) = \sup\{|\mathcal{A}| : \mathcal{A} \subset \{gA\}_{g \in G} \text{ is } |G|\text{-almost disjoint}\}.$$

A family $\mathcal{A}$ of sets is called κ -almost disjoint for a cardinal κ if $|A \cap A'| < \kappa$ for any distinct sets $A, A' \in \mathcal{A}$. So, being 1-almost disjoint is equivalent to being disjoint.

A subset $A \subset G$ with small packing index can be thought as large in a geometric sense because in this case the group G does not contain many disjoint translation copies of A . It is natural to compare this largeness property with other largeness properties that have topological or measure-theoretic nature. It turns out that a set of a group A can have small packing index (so can be large in geometric sense) and simultaneously be small in other senses. In [3] it was proved that each uncountable Polish Abelian group G contains a closed subset $A \subset G$ that has large packing index $\text{Pack}(A) = 1$ but is nowhere dense and Haar null in G . According to Theorem 16.3 [9], under CH (the Continuum Hypothesis), each Polish group G contains a subset A with packing index $\text{pack}(A) = 1$, which is *universally null* in the sense that A has measure zero with respect to any atomless Borel probability measure on G . In this paper we move further in this direction and prove that under

1991 *Mathematics Subject Classification.* 03E15; 03E50; 22A05; 54H05; 54H11.

Key words and phrases. Universally small set, universally meager set, universally null set, packing index, Polish group, coanalytic set.

CH each uncountable Abelian Polish group G contains a subset $A \subset G$ with large packing index $\text{Pack}(A) = 1$, which is *universally small* in the sense that it belongs to any ccc Borel σ -ideal on G . This fact combined with Theorem 6.3 of [3] allows us to construct universally small subsets of a given packing index in uncountable Polish Abelian groups.

Following Zakrzewski [11] we call a subset A of a Polish space X *universally small* if A belongs to each ccc σ -ideal with Borel base on X . By an *ideal* on a set X we understand a family $\mathcal{I}$ of subsets of X such that

- $\cup \mathcal{I} = X \notin \mathcal{I}$;
- for any sets $A, B \in \mathcal{I}$ we get $A \cup B \in \mathcal{I}$;
- for any sets $A \in \mathcal{I}$ and $B \subset X$ we get $A \cap B \in \mathcal{I}$.

An ideal $\mathcal{I}$ on a Polish space X is called

- a σ -*ideal* if $\cup \mathcal{A} \in \mathcal{I}$ for any countable subfamily $\mathcal{A} \subset \mathcal{I}$;
- an *ideal with Borel base* if each set $A \in \mathcal{I}$ lies in a Borel subset $B \in \mathcal{I}$;
- a *ccc ideal* if X contains no uncountable disjoint family of Borel subsets outside $\mathcal{I}$.

Standard examples of ccc Borel σ -ideals are the ideal $\mathcal{M}$ of meager subsets of a Polish space X and the ideal $\mathcal{N}$ of null subsets with respect to an atomless Borel σ -additive measure on X . This implies that a universally small subset A is universally null and universally meager. Following [10] we call a subset A of a Polish space X to be *universally meager* if for any Borel isomorphism $f : A \rightarrow 2^\omega$ the image $f(A)$ is meager in the Cantor cube 2^ω . Universally small sets were introduced by P. Zakrzewski [11] who constructed an uncountable universally small subset in each uncountable Polish space. It should be mentioned that there are models of ZFC [8, §5] in which all universally small sets in Polish spaces have cardinality $\leq \aleph_1 < \mathfrak{c}$. In such models any universally small set A in the real line has maximal possible packing index $\text{pack}(A) = \text{Pack}(A) = \mathfrak{c}$. This fact shows that the following theorem, which is the main result of this paper, necessarily has consistency nature and cannot be proved in ZFC.

Theorem 1. *Under CH, each uncountable Abelian Polish group G contains a universally small subset $A_0 \subset G$ with Packing index $\text{Pack}(A_0) = 1$.*

Combining this theorem with Theorem 6.4 of [3] we get

Corollary 1. *Under CH, for any cardinal $\kappa \in [2, \mathfrak{c}^+]$ with $\kappa \notin \{3, 4\}$ any uncountable Polish group G contains a universally small subset A with sharp packing index $\text{pack}^\sharp(A) = \kappa$.*

2. UNIVERSALLY SMALL SETS FROM COANALYTIC RANKS

In this section we describe a (known) method of constructing universally small sets, based on coanalytic ranks. Let us recall that a subset A of a Polish space X is

- *analytic* if A is the continuous image of a Polish space;
- *coanalytic* if $X \setminus A$ is analytic.

By Souslin's Theorem [5, 14.11], a subset of a Polish space is Borel if and only if it is analytic and coanalytic.

It is known [5, 34.4] that each coanalytic subset K of a Polish space X admits a *rank function* $\text{rank} : K \rightarrow \omega_1$ that has the following properties:

- (1) for every countable ordinal α the set $B_\alpha = \{x \in K : \text{rank}(x) \leq \alpha\}$ is Borel in X ;
- (2) each analytic subspace $A \subset K$ lies in some set B_α , $\alpha < \omega_1$.

The following fact is known and belongs to mathematical folklore (cf. [8, 5.3]). For the convenience of the reader we attach a short proof.

Lemma 1. *Let K be a coanalytic non-analytic set in a Polish space X and $\text{rank} : K \rightarrow \omega_1$ be a rank function. For any transfinite sequence of points $x_\alpha \in K \setminus B_\alpha$, $\alpha \in \omega_1$, the set $\{x_\alpha\}_{\alpha \in \omega_1}$ is universally small in X .*

Proof. Given any ccc Borel σ -ideal $\mathcal{I}$ on X , use the classical Szpilrajn-Marczewski Theorem [6, §11] to conclude that the coanalytic set K belongs to the completion $\mathcal{B}_{\mathcal{I}}(X) = \{A \subset X : \exists B \in \mathcal{B}(X) \ A \Delta B \in \mathcal{I}\}$ of the σ -algebra of Borel subsets of X by the ideal $\mathcal{I}$. Consequently, there is a Borel subset $B \subset K$ of X such that $K \setminus B \in \mathcal{I}$. By the property of the rank function, the Borel set B lies in B_β for some countable ordinal β . Then the set $\{x_\alpha\}_{\alpha < \omega_1}$ belongs to the σ -ideal $\mathcal{I}$, being the union of the countable set $\{x_\alpha\}_{\alpha \leq \beta}$ and the set $\{x_\alpha\}_{\beta < \alpha < \omega_1} \subset K \setminus B_\alpha \subset K \setminus B$ from the ideal $\mathcal{I}$. $\square$

In order to prove Theorem 1 we shall combine Lemma 1 with the following technical lemma that will be proved in Section 4.

Lemma 2. *For any uncountable Polish Abelian group G there are a non-empty open set $U \subset G$ and a co-analytic subset K of G such that $U \subset (K \setminus A) - (K \setminus A)$ for any analytic subspace $A \subset K$ of G .*

3. PROOF OF THEOREM 1

Assume the Continuum Hypothesis. Given an uncountable Polish Abelian group G we need to construct a universally small subset $A \subset G$ with $\text{Pack}(A) = 1$. We shall use the additive notation for denoting the group operation on G . So, 0 will denote the neutral element of G . For two subsets $A, B \subset G$ we put $A + B = \{a + b : a \in A, b \in B\}$ and $A - B = \{a - b : a \in A, b \in B\}$.

By Lemma 2, there are a non-empty open set $U \subset G$ and a coanalytic subset K such that $U \subset (K \setminus B) - (K \setminus B)$ for any Borel subset $B \subset K$ of G . This implies that the coanalytic set K is not Borel in G . Let $\text{rank} : K \rightarrow \omega_1$ be a rank function for K . This function induces the decomposition $K = \bigcup_{\alpha < \omega_1} B_\alpha$ into Borel sets $B_\alpha = \{x \in K : \text{rank}(x) \leq \alpha\}$, $\alpha < \omega_1$, such that each Borel subset $B \subset K$ of G lies in some set B_α , $\alpha < \omega_1$.

The Continuum Hypothesis allows us to choose an enumeration $U = \{u_\alpha\}_{\alpha < \omega_1}$ of the open set U such that for every $u \in U$ the set $\Omega_u = \{\alpha < \omega_1 : u_\alpha = u\}$ is uncountable. The separability of G yields a countable subset $C \subset G$ such that $G = C + U$.

By induction, for every $\alpha < \omega_1$ find two points $x_\alpha, y_\alpha \in K \setminus (B_\alpha \cup \{x_\beta : \beta < \alpha\})$ such that $x_\alpha - y_\alpha = u_\alpha$. Such a choice is always possible as $U \subset (K \setminus B) - (K \setminus B)$ for any Borel subset $B \subset K$ of G . Lemma 1 guarantees that the sets $\{x_\alpha\}_{\alpha < \omega_1}$ and $\{y_\alpha\}_{\alpha < \omega_1}$ are universally small in G and so is the set $A = \{c + x_\alpha, y_\alpha : c \in C, \alpha < \omega_1\}$. It remains to prove that $\text{Pack}(A) = 1$. This equality will follow as soon as we check that for every point $z \in G$ the intersection $A \cap (z + A)$ has cardinality of continuum. Since $C + U = G$, we can find elements $c \in C$ and $u \in U$ such that $z = c + u$. The choice of the enumeration $\{u_\alpha\}_{\alpha < \omega_1}$ guarantees that the set $\Omega_u = \{\alpha < \omega_1 : u_\alpha = u\}$ has cardinality continuum. Now observe that for every $\alpha \in \Omega_u$ we get $z = c + u = c + u_\alpha = c + x_\alpha - y_\alpha$ and hence $c + x_\alpha = z + y_\alpha \in A \cap (z + A)$, which implies that the intersection $A \cap (z + A) \supset \{c + x_\alpha\}_{\alpha \in \Omega_u}$ has cardinality of continuum.

4. PROOF OF LEMMA 2

Fix an invariant metric $d \leq 1$ generating the topology of G . This metric is complete because the group G is Polish. The metric d induces a norm $\|\cdot\| : G \rightarrow [0, 1]$ on G defined by $\|x\| = d(x, 0)$. For

an $\varepsilon > 0$ by $B(\varepsilon) = \{x \in G : \|x\| < \varepsilon\}$ and $\bar{B}(\varepsilon) = \{x \in G : \|x\| \leq \varepsilon\}$ we shall denote the open and closed ε -balls centered at zero.

We define a subset D of G to be ε -separated if $d(x, y) \geq \varepsilon$ for any distinct points $x, y \in D$. By Zorn's Lemma, each ε -separated subset S of any subset $A \subset G$ can be enlarged to a maximal ε -separated subset $\tilde{S}$ of A . This set $\tilde{S}$ is ε -net for A in the sense that for each point $a \in A$ there is a point $s \in \tilde{S}$ with $d(a, s) < \varepsilon$.

Fix any non-zero element $a_{-1} \in G$ and let $\varepsilon_{-1} = \frac{1}{12}\|a_{-1}\|$. By induction we can define a sequence $(\varepsilon_n)_{n \in \omega}$ of positive real numbers and a sequence $(a_n)_{n \in \omega}$ of points of the group G such that

- $16\varepsilon_n \leq \|a_n\| < \varepsilon_{n-1}$ for every $n \in \omega$.

For every $n \in \omega$, fix a maximal $2\varepsilon_n$ -separated subset $X_n \ni 0$ in the ball $B(2\varepsilon_{n-1})$.

The choice of the sequence (ε_n) guarantees that the series $\sum_{n \in \omega} \varepsilon_n$ is convergent and thus for any sequence $(x_n)_{n \in \omega} \in \prod_{n \in \omega} X_n$ the series $\sum_{n \in \omega} x_n$ is convergent in G (because $\|x_n\| < 2\varepsilon_{n-1}$ for all $n \in \mathbb{N}$). Therefore the following subsets of the group G are well-defined:

$$\begin{aligned}\Sigma_0 &= \left\{ \sum_{n \in \omega} x_{2n} : (x_{2n})_{n \in \omega} \in \prod_{n \in \omega} X_{2n} \right\}, \\ \Sigma_1 &= \left\{ \sum_{n \in \omega} x_{2n+1} : (x_{2n+1})_{n \in \omega} \in \prod_{n \in \omega} X_{2n+1} \right\}.\end{aligned}$$

These sets have the following properties:

Claim 1. (1) $\Sigma_0 \cup \Sigma_1 \subset B(4\varepsilon_{-1})$.

(2) $B(2\varepsilon_{-1}) \subset \Sigma_1 + \Sigma_0$.

(3) For every $i \in \{0, 1\}$ the closure $\overline{\Sigma_i - \Sigma_i}$ of the set $\Sigma_i - \Sigma_i$ in G is not a neighborhood of zero.

Proof. 1. For every point $x \in \Sigma_0 \cup \Sigma_1$ we can find a sequence $(x_n)_{n \in \omega} \in \prod_{n \in \omega} X_n$ with $x = \sum_{n=0}^{\infty} x_n$ and observe that

$$\|x\| \leq \sum_{n=0}^{\infty} \|x_n\| \leq \sum_{n=0}^{\infty} 2\varepsilon_{n-1} < \sum_{n \in \omega} \frac{2\varepsilon_{-1}}{16^n} < 4\varepsilon_{-1}.$$

2. Given any point $x \in B(2\varepsilon_{-1})$, find a point $x_0 \in X_0$ such that $\|x - x_0\| < 2\varepsilon_0$. Such a point x_0 exists as the set X_0 is a $2\varepsilon_0$ -net in $B(2\varepsilon_{-1})$. Continuing by induction, for every $n \in \omega$ find a point $x_n \in X_n$ such that $\|x - \sum_{i=0}^n x_i\| < 2\varepsilon_n$. After completing the inductive construction, we obtain a sequence $(x_n)_{n \in \omega} \in \prod_{n \in \omega} X_n$ such that

$$x = \sum_{n \in \omega} x_n = \sum_{n \in \omega} x_{2n} + \sum_{n \in \omega} x_{2n+1} \in \Sigma_0 + \Sigma_1.$$

3. We shall give a detail proof of the third statement for $i = 0$ (for $i = 1$ the proof is analogous). Since the sequence $(a_{2k+1})_{k \in \omega}$ converges to zero, it suffices to show that $d(a_{2k+1}, \Sigma_0 - \Sigma_0) > 0$ for all $k \in \omega$.

Given two points $x, y \in \Sigma_0$, we shall prove that $d(a_{2k+1}, x - y) \geq \varepsilon_{2k+1}$. If $x = y$, then $d(a_{2k+1}, x - y) = d(a_{2k+1}, 0) = \|a_{2k+1}\| > \varepsilon_{2k+1}$ by the choice of a_{2k+1} . So, we assume that $x \neq y$. Find infinite sequences $(x_{2n})_{n \in \omega}, (y_{2n})_{n \in \omega} \in \prod_{n \in \omega} X_{2n}$ with $x = \sum_{n \in \omega} x_{2n}$ and $y = \sum_{n \in \omega} y_{2n}$.

Let $m = \min\{n \in \omega : x_{2n} \neq y_{2n}\}$. If $m \geq k + 1$, then

$$\begin{aligned}\|x - y\| &= \left\| \sum_{n \geq m} x_{2n} - y_{2n} \right\| \leq \sum_{n \geq m} \|x_{2n}\| + \|y_{2n}\| \leq \\ &\leq 2 \sum_{n \geq m} 2\varepsilon_{2n-1} \leq 8\varepsilon_{2m-1} \leq 8\varepsilon_{2k+1} < \|a_{2k+1}\| - \varepsilon_{2k+1}\end{aligned}$$

and hence $d(x - y, a_{2k+1}) \geq \varepsilon_{2k+1}$.

If $m \leq k$, then

$$\begin{aligned} \|x - y\| &= \|(x_{2m} - y_{2m}) + \sum_{n>m} (x_{2n} - y_{2n})\| \geq \|x_{2m} - y_{2m}\| - \sum_{n>m} (\|x_{2n}\| + \|y_{2n}\|) \geq \\ &\geq 2\varepsilon_{2m} - 2 \sum_{n>m} 2\varepsilon_{2n-1} \geq 2\varepsilon_{2m} - 8\varepsilon_{2m+1} \geq \frac{3}{2}\varepsilon_{2m} \geq \frac{3}{2}\varepsilon_{2k} > \|a_{2k+1}\| + \frac{1}{2}\varepsilon_{2k} \end{aligned}$$

according to the choice of the point a_{2k+1} . Consequently, $d(x - y, a_{2k+1}) \geq \frac{1}{2}\varepsilon_{2k} \geq \varepsilon_{2k+1}$. $\square$

A subset C of G will be called a *Cantor set* in G if C is homeomorphic to the Cantor cube $\{0, 1\}^\omega$. By the classical Brouwer's Theorem [5, 7.4], this happens if and only if C is compact, zero-dimensional and has no isolated points.

Claim 2. *For every $i \in \{0, 1\}$ there is a Cantor set $C_i \subset B(\varepsilon_0)$ such that the map $h_i : C_i \times \overline{\Sigma}_i \rightarrow G$, $h_i : (x, y) \mapsto x + y$, is a closed topological embedding.*

Proof. Taking into account that $\overline{\Sigma}_i - \overline{\Sigma}_i = \overline{\Sigma}_i - \overline{\Sigma}_i$ is not a neighborhood of zero in G , and repeating the proof of Lemma 2.1 of [3], we can construct a Cantor set $C_i \subset B(\varepsilon_0)$ such that for any distinct points $x, y \in C_i$ the shifts $x + \overline{\Sigma}_i$ and $y + \overline{\Sigma}_i$ are disjoint. This implies that the map $h_i : C_i \times \overline{\Sigma}_i \rightarrow G$, $h_i : (x, y) \mapsto x + y$, is injective. Taking into account that the set C_i is compact and $\overline{\Sigma}_i$ is closed in G , one can check that the map h_i is closed and hence a closed topological embedding. $\square$

Observe that for every $i \in \{0, 1\}$ the embedding h_i has image $h_i(C_i \times \overline{\Sigma}_i) = C_i + \overline{\Sigma}_i \subset B(\varepsilon_0) + \overline{B}(4\varepsilon_{-1}) \subset B(5\varepsilon_{-1})$. Now we modify the closed embeddings h_0 and h_1 to closed embeddings

$$\tilde{h}_0 : C_0 \times \overline{\Sigma}_0 \rightarrow G, \quad \tilde{h}_0 : (x, y) \mapsto a_{-1} + x + y$$

and

$$\tilde{h}_1 : C_1 \times \overline{\Sigma}_1 \rightarrow G, \quad \tilde{h}_1(x, y) = -x - y.$$

These embeddings have images $\tilde{h}_0(C_0 \times \overline{\Sigma}_0) \subset a_{-1} + B(5\varepsilon_{-1})$ and $\tilde{h}_1(C_1 \times \overline{\Sigma}_1) \subset a_{-1} - B(5\varepsilon_{-1}) = a_{-1} + B(5\varepsilon_{-1})$. Since $\|a_{-1}\| = 12\varepsilon_{-1}$, we conclude that the closed subsets $\tilde{h}_i(C_i \times \overline{\Sigma}_i)$, $i \in \{0, 1\}$, of G are disjoint.

For every $i \in \{0, 1\}$ fix a coanalytic non-analytic subset K_i in the Cantor set C_i . It follows that the disjoint union $K = \bigcup_{i=0}^1 \tilde{h}_i(K_i \times \overline{\Sigma}_i)$ is a coanalytic subset of G .

The following claim completes the proof of the lemma and shows that the coanalytic set K and the open set $U = a_{-1} + B(\varepsilon_{-1})$ have the required property.

Claim 3. $U \subset (K \setminus A) - (K \setminus A)$ for any analytic subspace $A \subset K$.

Proof. Given an analytic subspace $A \subset K$, for every $i \in \{0, 1\}$, consider its preimage $A_i = \tilde{h}_i^{-1}(A) \subset C_i \times \overline{\Sigma}_i$ and its projection $\text{pr}_i(A_i)$ onto the Cantor set C_i . It follows from $A \subset K$ and $\bigcap_{i=0}^1 \tilde{h}_i(C_i \times \overline{\Sigma}_i) = \emptyset$ that each set A_i is an analytic subspace of the coanalytic set K_i . Since the space K_i is not analytic, there is a point $c_i \in K_i \setminus \text{pr}_i(A_i)$. It follows that

$$\bigcup_{i=0}^1 \tilde{h}_i(\{c_i\} \times \Sigma_i) = (a_{-1} + c_0 + \Sigma_0) \cup (-c_1 - \Sigma_1) \subset K \setminus A$$

and hence

$$(K \setminus A) - (K \setminus A) \supset a_{-1} + c_0 + \Sigma_0 + c_1 + \Sigma_1 \supset a_{-1} + c_0 + c_1 + B(2\varepsilon_{-1}) \supset a_{-1} + B(\varepsilon_{-1}) = U$$

according to Claim 1(2). The inclusion $B(\varepsilon_{-1}) \subset c_0 + c_1 + B(2\varepsilon_{-1})$ follows from $c_0 + c_1 \in C_0 + C_1 \subset B(\varepsilon_0) + B(\varepsilon_0) \subset B(2\varepsilon_0) \subset B(\varepsilon_{-1})$. $\square$

REFERENCES

- [1] T. Banakh, N. Lyaskovska, *Weakly P -small not P -small subsets in Abelian groups*, Algebra and Discr. Math. N.3 (2006), 29–34.
- [2] T. Banakh, N. Lyaskovska, *Weakly P -small not P -small subsets in groups*, J. Algebra Comput. **18**:1 (2008), 1–6
- [3] T. Banakh, N. Lyaskovska, D. Repovs, *Packing index of subsets in Polish groups*, Notre Dame J. Formal Logic. **50**:4 (2009), 453–468.
- [4] D. Dikranjan, I. Protasov, *Every infinite group can be generated by P -small subset*, Appl. Gen. Topology, **7** (2006), 265–268.
- [5] A. Kechris, *Classical Descriptive Set Theory*, Springer-Verlag, New York, 1995.
- [6] K. Kuratowski, *Topology, I*, Mir, Moskov, 1966.
- [7] N. Lyaskovska, *Constructing subsets of a given packing index in abelian groups*, Acta Univ. Carolin. Math. Phys. **48**:2 (2007), 69–80.
- [8] A.W. Miller, *Special subsets of the real line*, in: Handbook of set-theoretic topology, North-Holland, Amsterdam, (1984), 201–233.
- [9] I. Protasov, T. Banakh, *Ball structures and coloring of graphs and groups*, VNTL Publ., Lviv, 2003.
- [10] P. Zakrzewski, *Universally meager sets*, Proc. Amer. Math. Soc. **129**:6 (2001), 1793–1798.
- [11] P. Zakrzewski, *On a construction of universally small sets*, Real Anal. Exchange **28**:1 (2002/03), 221–226.

INSTYTUT MATEMATYKI, UNIWERSYTET JANA KOCHANOWSKIEGO, KIELCE, POLAND

DEPARTMENT OF MATHEMATICS, IVAN FRANKO NATIONAL UNIVERSITY OF LVIV, UKRAINE

E-mail address: t.o.banakh@gmail.com, lyaskovska@yahoo.com